



Stable inhibitors of the 5-lipoxygenase pathway for treating asthma and cancer

IDEA

Introduction of metabolically stable carboranes into inhibitors of the 5-LO pathway to enhance the pharmacokinetic profile of related drugs.

DEMONSTRATOR

Introduction of carboranes maintains the inhibitory potential for 5-LO, but the cytotoxicity for invasive melanoma and colon cancer cell lines is strongly increased. IC_{50} = 10 -20 μ molar

PROTOTYPE

Mice with colon adenocarcinoma show an improved response to the investigated compounds (clear reduction of tumor). No acute toxicity.

BACKGROUND/ MEDICAL PROBLEM

5-Lipoxygenase (5-LO) is an enzyme that catalyses the conversion of arachidonic acid to biologically active leukotrienes (LTs). LTs are cellular signalling mediators involved in inflammatory diseases, such as asthma, atherosclerosis, cardiovascular diseases and diverse types of cancer, e.g. acute myeloid leukaemia. Competitive LT receptor antagonists such as *montelukast* and *zafirlukast* have been developed to decrease the symptoms of an overexpression of LTs. They bind to the cysteinyl leukotriene receptor ($CysLT_1$), which is disadvantageous, since lipoxygenases also produce other LTs that bind to other receptors. Therefore, the inhibition of 5-LO is beneficial, since the production of all LTs is decreased. *Zileuton* for treatment of asthma is the only drug on the market inhibiting the 5-LO pathway. However, the pharmacological profile of *zileuton* is disadvantageous, since it is rapidly metabolised.

POTENTIAL APPLICATION

- Treatment of 5-LO dependent tumors
- Tumor imaging
- Boron neutron capture therapy (BNCT)


ADVANTAGES

- 5-Lipoxygenase is a therapeutic target involved in diseases like asthma, atherosclerosis and cancer
- Carboranes improve pharmacokinetic behavior
- Carborane-based 5-LO inhibitors increase the selective cytotoxicity for cancer cells
- Easy and fast synthesis in three steps

Project 17008

TECHNOLOGY
OFFER

Carborane-based 5-LO inhibitors



TECHNOLOGY/SOLUTION

We have investigated the substitution of metabolisable phenyl rings of lipoxygenase inhibitors by metabolically stable boron compounds, namely, carboranes ($C_2B_{10}H_{12}$). Carboranes are icosahedral boron clusters in which two boron atoms are replaced by carbon atoms. They are non-toxic, have remarkable biological stability, and the carbon atoms can be functionalised easily and selectively. We have found that the introduction of carboranes does not influence the inhibitory potential for the 5-LO pathway, but strongly and selectively increases the cytotoxicity for cancer cells.

Figure

Left: Leukotrienes, produced by 5-lipoxygenase, lead to angiogenesis and metastasis of cancer cells. **Right:** Introduction of carboranes leads to highly selective and metabolically stable lipoxygenase inhibitors which can prevent angiogenesis.

FURTHER READING:

J. Roos et al., *Pharmacol. Ther.*, **2016**, 157, 43–64 ,
(<https://doi.org/10.1016/j.pharmthera.2015.11.001>)

O. Werz, *Planta Med.*, **2007**, 73, 1331–1357. (DOI: [10.1055/s-2007-990242](https://doi.org/10.1055/s-2007-990242))

STATUS OF PROPRIETARY RIGHT

Patent is pending

COOPERATION OPTIONS

- R&D Agreement
- License Agreement
- Ownership Agreement

CONTACT

LEIPZIG UNIVERSITY
Life Science Transfer Office
Deutscher Platz 5
04103 Leipzig
Germany

technologietransfer@bbz.uni-leipzig.de
Tel.: +49 341 97-31387

RESPONSIBLE RESEARCH INSTITUTE

LEIPZIG UNIVERSITY
Institute of Chemistry/
Institute of Biochemistry
<https://research.uni-leipzig.de/hh>
<https://biochemie.lw.uni-leipzig.de/arbeitsgruppen/biochemie-und-bioorganische-chemie>